

**q -ANALOGUE OF p -ADIC $\log \Gamma$ TYPE FUNCTIONS
ASSOCIATED WITH MODIFIED q -EXTENSION OF GENOCCHI
NUMBERS WITH WEIGHT α AND β**

SERKAN ARACI AND MEHMET ACIKGOZ

ABSTRACT. The fundamental aim of this paper is to describe q -Analogue of p -adic log gamma functions with weight alpha and beta. Moreover, we give relationship between p -adic q -log gamma functions with weight (α, β) and q -extension of Genocchi numbers with weight alpha and beta and modified q -Euler numbers with weight α

1. INTRODUCTION

Assume that p be a fixed odd prime number. Throughout this paper \mathbb{Z} , \mathbb{Z}_p , \mathbb{Q}_p and \mathbb{C}_p will denote by the ring of integers, the field of p -adic rational numbers and the completion of the algebraic closure of \mathbb{Q}_p , respectively. Also we denote $\mathbb{N}^* = \mathbb{N} \cup \{0\}$ and $\exp(x) = e^x$. Let $v_p : \mathbb{C}_p \rightarrow \mathbb{Q} \cup \{\infty\}$ (\mathbb{Q} is the field of rational numbers) denote the p -adic valuation of \mathbb{C}_p normalized so that $v_p(p) = 1$. The absolute value on \mathbb{C}_p will be denoted as $|\cdot|_p$, and $|x|_p = p^{-v_p(x)}$ for $x \in \mathbb{C}_p$. When one talks of q -extensions, q is considered in many ways, e.g. as an indeterminate, a complex number $q \in \mathbb{C}$, or a p -adic number $q \in \mathbb{C}_p$. If $q \in \mathbb{C}$ we assume that $|q| < 1$. If $q \in \mathbb{C}_p$, we assume $|1 - q|_p < p^{-\frac{1}{p-1}}$, so that $q^x = \exp(x \log q)$ for $|x|_p \leq 1$. We use the following notation

$$(1.1) \quad [x]_q = \frac{1 - q^x}{1 - q}, \quad [x]_{-q} = \frac{1 - (-q)^x}{1 + q}$$

where $\lim_{q \rightarrow 1} [x]_q = x$; cf. [1-24].

For a fixed positive integer d with $(d, p) = 1$, we set

$$\begin{aligned} X &= X_d = \varprojlim_{\mathbb{N}} \mathbb{Z}/dp^N \mathbb{Z}, \\ X^* &= \bigcup_{\substack{0 < a < dp \\ (a, p) = 1}} a + dp\mathbb{Z}_p \end{aligned}$$

and

$$a + dp^N \mathbb{Z}_p = \{x \in X \mid x \equiv a \pmod{dp^N}\},$$

where $a \in \mathbb{Z}$ satisfies the condition $0 \leq a < dp^N$.

It is known that

$$\mu_q(x + p^N \mathbb{Z}_p) = \frac{q^x}{[p^N]_q}$$

Date: January 20, 2012.

2000 Mathematics Subject Classification. Primary 46A15, Secondary 41A65.

Key words and phrases. Modified q -Genocchi numbers with weight alpha and beta, Modified q -Euler numbers with weight alpha and beta, p -adic log gamma functions.

is a distribution on X for $q \in \mathbb{C}_p$ with $|1 - q|_p \leq 1$.

Let $UD(\mathbb{Z}_p)$ be the set of uniformly differentiable function on \mathbb{Z}_p . We say that f is a uniformly differentiable function at a point $a \in \mathbb{Z}_p$, if the difference quotient

$$F_f(x, y) = \frac{f(x) - f(y)}{x - y}$$

has a limit $f'(a)$ as $(x, y) \rightarrow (a, a)$ and denote this by $f \in UD(\mathbb{Z}_p)$. The p -adic q -integral of the function $f \in UD(\mathbb{Z}_p)$ is defined by

$$(1.2) \quad I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x) q^x$$

The bosonic integral is considered by Kim as the bosonic limit $q \rightarrow 1$, $I_1(f) = \lim_{q \rightarrow 1} I_q(f)$. Similarly, the p -adic fermionic integration on \mathbb{Z}_p defined by Kim as follows:

$$I_{-q}(f) = \lim_{q \rightarrow -q} I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x)$$

Let $q \rightarrow 1$, then we have p -adic fermionic integral on \mathbb{Z}_p as follows:

$$I_{-1}(f) = \lim_{q \rightarrow -1} I_q(f) = \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) (-1)^x.$$

Stirling asymptotic series are defined by

$$(1.3) \quad \log\left(\frac{\Gamma(x+1)}{\sqrt{2\pi}}\right) = \left(x - \frac{1}{2}\right) \log x + \sum_{n=1}^{\infty} \frac{(-1)^{n+1} B_{n+1}}{n(n+1)} \frac{1}{x^n} - x$$

where B_n are familiar n -th Bernoulli numbers cf. [6, 8, 9, 25].

Recently, Araci et al. defined modified q -Genocchi numbers and polynomials with weight α and β in [4, 5] by the means of generating function:

$$(1.4) \quad \sum_{n=0}^{\infty} g_{n,q}^{(\alpha,\beta)}(x) \frac{t^n}{n!} = t \int_{\mathbb{Z}_p} q^{-\beta\xi} e^{[x+\xi]_{q^\alpha} t} d\mu_{-q^\beta}(\xi)$$

So from above, we easily get Witt's formula of modified q -Genocchi numbers and polynomials with weight α and β as follows:

$$(1.5) \quad \frac{g_{n+1,q}^{(\alpha,\beta)}(x)}{n+1} = \int_{\mathbb{Z}_p} q^{-\beta\xi} [x+\xi]_{q^\alpha}^n d\mu_{-q^\beta}(\xi)$$

where $g_{n,q}^{(\alpha,\beta)}(0) := g_{n,q}^{(\alpha,\beta)}$ are modified q extension of Genocchi numbers with weight α and β cf. [4,5].

In [21], Rim and Jeong are defined modified q -Euler numbers with weight α as follows:

$$(1.6) \quad \tilde{\xi}_{n,q}^{(\alpha)} = \int_{\mathbb{Z}_p} q^{-t} [t]_{q^\alpha} d\mu_{-q}(t)$$

From expressions of (1.5) and (1.6), we get the following Proposition 1:

Proposition 1. *The following*

$$(1.7) \quad \tilde{\xi}_{n,q}^{(\alpha)} = \frac{g_{n+1,q}^{(\alpha,1)}}{n+1}$$

is true.

In previous paper [6], Araci, Acikgoz and Park introduced weighted q -Analogue of p -Adic log gamma type functions and they derived some interesting identities in Analytic Numbers Theory and in p -Adic Analysis. They were motivated from paper of T. Kim by "On a q -analogue of the p -adic log gamma functions and related integrals, *J. Number Theory*, 76 (1999), no. 2, 320-329." We also introduce q -Analogue of p -Adic log gamma type function with weight α and β . We derive in this paper some interesting identities this type of functions.

On p -adic $\log \Gamma$ function with weight α and β

In this part, from (1.2), we begin with the following nice identity:

$$(1.8) \quad I_{-q}^{(\beta)}(q^{-\beta x} f_n) + (-1)^{n-1} I_{-q}^{(\beta)}(q^{-\beta x} f) = [2]_{q^\beta} \sum_{l=0}^{n-1} (-1)^{n-1-l} f(l)$$

where $f_n(x) = f(x+n)$ and $n \in \mathbb{N}$ (see [4]).

In particular for $n = 1$ into (1.8), we easily see that

$$(1.9) \quad I_{-q}^{(\beta)}(q^{-\beta x} f_1) + I_{-q}^{(\beta)}(q^{-\beta x} f) = [2]_{q^\beta} f(0).$$

With the simple application, it is easy to indicate as follows:

$$(1.10) \quad ((1+x) \log(1+x))' = 1 + \log(1+x) = 1 + \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n(n+1)} x^n$$

where $((1+x) \log(1+x))' = \frac{d}{dx}((1+x) \log(1+x))$

By expression of (1.10), we can derive

$$(1.11) \quad (1+x) \log(1+x) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n(n+1)} x^{n+1} + x + c, \text{ where } c \text{ is constant.}$$

If we take $x = 0$, so we get $c = 0$. By expression of (1.10) and (1.11), we easily see that,

$$(1.12) \quad (1+x) \log(1+x) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n(n+1)} x^{n+1} + x.$$

It is considered by T. Kim for q -analogue of p adic locally analytic function on $\mathbb{C}_p \setminus \mathbb{Z}_p$ as follows:

$$(1.13) \quad G_{p,q}(x) = \int_{\mathbb{Z}_p} [x+\xi]_q \left(\log [x+\xi]_q - 1 \right) d\mu_{-q}(\xi) \text{ (for detail, see[5,6]).}$$

By the same motivation of (1.13), in previous paper [6], q -analogue of p -adic locally analytic function on $\mathbb{C}_p \setminus \mathbb{Z}_p$ with weight α is considered

$$(1.14) \quad G_{p,q}^{(\alpha)}(x) = \int_{\mathbb{Z}_p} [x+\xi]_{q^\alpha} \left(\log [x+\xi]_{q^\alpha} - 1 \right) d\mu_{-q}(\xi)$$

In particular $\alpha = 1$ into (1.14), we easily see that, $G_{p,q}^{(1)}(x) = G_{p,q}(x)$.

With the same manner, we introduce q -Analogue of p -adic locally analytic function on $\mathbb{C}_p \setminus \mathbb{Z}_p$ with weight α and β as follows:

$$(1.15) \quad G_{p,q}^{(\alpha,\beta)}(x) = \int_{\mathbb{Z}_p} q^{-\beta\xi} [x + \xi]_{q^\alpha} \left(\log [x + \xi]_{q^\alpha} - 1 \right) d\mu_{-q^\beta}(\xi)$$

From expressions of (1.9) and (1.16), we state the following Theorem:

Theorem 1. *The following identity holds:*

$$G_{p,q}^{(\alpha,\beta)}(x+1) + G_{p,q}^{(\alpha,\beta)}(x) = [2]_{q^\beta} [x]_{q^\alpha} \left(\log [x]_{q^\alpha} - 1 \right).$$

It is easy to show that,

$$(1.16) \quad \begin{aligned} [x + \xi]_{q^\alpha} &= \frac{1 - q^{\alpha(x+\xi)}}{1 - q^\alpha} \\ &= \frac{1 - q^{\alpha x} + q^{\alpha x} - q^{\alpha(x+\xi)}}{1 - q^\alpha} \\ &= \left(\frac{1 - q^{\alpha x}}{1 - q^\alpha} \right) + q^{\alpha x} \left(\frac{1 - q^{\alpha\xi}}{1 - q^\alpha} \right) \\ &= [x]_{q^\alpha} + q^{\alpha x} [\xi]_{q^\alpha} \end{aligned}$$

Substituting $x \rightarrow \frac{q^{\alpha x} [\xi]_{q^\alpha}}{[x]_{q^\alpha}}$ into (1.12) and by using (1.16), we get interesting formula:

$$(1.17) \quad [x + \xi]_{q^\alpha} \left(\log [x + \xi]_{q^\alpha} - 1 \right) = \left([x]_{q^\alpha} + q^{\alpha x} [\xi]_{q^\alpha} \right) \log [x]_{q^\alpha} + \sum_{n=1}^{\infty} \frac{(-q^{\alpha x})^{n+1}}{n(n+1)} \frac{[\xi]_{q^\alpha}^{n+1}}{[x]_{q^\alpha}^n} - [x]_{q^\alpha}$$

If we substitute $\alpha = 1$ into (1.17), we get Kim's q -Analogue of p -adic log gamma fuction (for detail, see[8]).

From expression of (1.2) and (1.17), we obtain worthwhile and interesting theorems as follows:

Theorem 2. *For $x \in \mathbb{C}_p \setminus \mathbb{Z}_p$ the following*

$$(1.18) \quad G_{p,q}^{(\alpha,\beta)}(x) = \left(\frac{[2]_{q^\beta}}{2} [x]_{q^\alpha} + q^{\alpha x} \frac{g_{2,q}^{(\alpha,\beta)}}{2} \right) \log [x]_{q^\alpha} + \sum_{n=1}^{\infty} \frac{(-q^{\alpha x})^{n+1}}{n(n+1)(n+2)} \frac{g_{n+1,q}^{(\alpha,\beta)}}{[x]_{q^\alpha}^n} - [x]_{q^\alpha} \frac{[2]_{q^\beta}}{2}$$

is true.

Corollary 1. *Taking $q \rightarrow 1$ into (1.18), we get nice identity:*

$$G_{p,1}^{(\alpha,\beta)}(x) = \left(x + \frac{G_2}{2} \right) \log x + \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n(n+1)(n+2)} \frac{G_{n+1}}{x} - x$$

where G_n are called famous Genocchi numbers.

Theorem 3. *The following nice identity*

$$(1.19) \quad G_{p,q}^{(\alpha,1)}(x) = \left(\frac{[2]_q}{2} [x]_{q^\alpha} + q^{\alpha x} \tilde{\xi}_{1,q}^{(\alpha)} \right) \log [x]_{q^\alpha} + \sum_{n=1}^{\infty} \frac{(-q^{\alpha x})^{n+1}}{n(n+1)} \frac{\tilde{\xi}_{n,q}^{(\alpha)}}{[x]_{q^\alpha}^n} - \frac{[2]_q}{2} [x]_{q^\alpha}$$

is true.

Corollary 2. *Putting $q \rightarrow 1$ into (1.19), we have the following identity:*

$$G_{p,1}^{(\alpha,\beta)}(x) = (x + E_1) \log x + \sum_{n=1}^{\infty} \frac{(-1)^{n+1} E_n}{n(n+1) x^n} - x$$

where E_n are familiar Euler numbers.

REFERENCES

- [1] Araci, S., Acikgoz, M., and Seo, J-J., A note on the weighted q -Genocchi numbers and polynomials with Their Interpolation Function, Accepted in Honam Mathematical Journal.
- [2] Araci, S., Erdal, D., and Seo., J-J., A study on the fermionic p adic q - integral representation on \mathbb{Z}_p associated with weighted q -Bernstein and q -Genocchi polynomials, Abstract and Applied Analysis, Volume **2011**, Article ID 649248, 10 pages.
- [3] Araci, S., Seo, J-J., and Erdal, D., New construction weighted (h,q) -Genocchi numbers and polynomials related to Zeta type function, Discrete Dynamics in nature and Society, Volume **2011**, Article ID 487490, 7 pages.
- [4] Araci, S., Acikgoz, M., Qi, Feng., and Jolany, H., A note on the modified q -Genocchi numbers and polynomials with weight (α, β) and Their Interpolation function at negative integer, submitted.
- [5] Araci, S., Acikgoz, M., and Ryoo, C-S., A note on the values of the weighted q -Bernstein Polynomials and Modified q -Genocchi Numbers with weight α and β via the p -adic q -integral on \mathbb{Z}_p , submitted
- [6] Araci, S., Acikgoz, M., Park, K-H., A note on the q -Analogue of Kim's p -adic \log gamma functions associated with q -extension of Genocchi and Euler polynomials with weight α , submitted.
- [7] Acikgoz, M., and Simsek, Y., On multiple interpolation functions of the Nörlund type q -Euler polynomials, Abstract and Applied Analysis, Volume **2009**, Article ID 382574, 14 pages.
- [8] Kim, T., A note on the q -analogue of p -adic \log gamma function, arXiv:0710.4981v1 [math.NT].
- [9] Kim, T., On a q -analogue of the p -adic \log gamma functions and related integrals, J. Number Theory, **76** (1999), no. 2, 320-329.
- [10] Kim, T., On the q -extension of Euler and Genocchi numbers, J. Math. Anal. Appl. 326 (2007) 1458-1465.
- [11] Kim, T., On the multiple q -Genocchi and Euler numbers, Russian J. Math. Phys. 15 (4) (2008) 481-486. arXiv:0801.0978v1 [math.NT]
- [12] Kim, T., On the weighted q -Bernoulli numbers and polynomials, Advanced Studies in Contemporary Mathematics 21(2011), no.2, p. 207-215, <http://arxiv.org/abs/1011.5305>.
- [13] Kim, T., q -Volkenborn integration, Russ. J. Math. phys. 9(2002) , 288-299.
- [14] Kim, T., An invariant p -adic q -integrals on \mathbb{Z}_p , Applied Mathematics Letters, vol. 21, pp. 105-108, 2008.
- [15] Kim, T., q -Euler numbers and polynomials associated with p -adic q -integrals, J. Nonlinear Math. Phys., 14 (2007), no. 1, 15-27.
- [16] Kim, T., New approach to q -Euler polynomials of higher order, Russ. J. Math. Phys., 17 (2010), no. 2, 218-225.
- [17] Kim, T., Some identities on the q -Euler polynomials of higher order and q -Stirling numbers by the fermionic p -adic integral on \mathbb{Z}_p , Russ. J. Math. Phys., 16 (2009), no.4, 484-491.
- [18] Kim, T. and Rim, S.-H., On the twisted q -Euler numbers and polynomials associated with basic q - l -functions, Journal of Mathematical Analysis and Applications, vol. 336, no. 1, pp. 738-744, 2007.
- [19] Kim, T., On p -adic q - l -functions and sums of powers, J. Math. Anal. Appl. (2006), doi:10.1016/j.jmaa.2006.07.071
- [20] Ryoo, C. S., A note on the weighted q -Euler numbers and polynomials, Advan. Stud. Contemp. Math. 21(2011), 47-54.
- [21] Rim, S-H., and Jeong, J., A note on the modified q -Euler numbers and Polynomials with weight α , International Mathematical Forum, Vol. 6, 2011, no. 65, 3245-3250.
- [22] Simsek, Y., Theorems on twisted L -function and twisted Bernoulli numbers, Advan. Stud. Contemp. Math., 11(2005), 205-218.

- [23] Simsek, Y., Twisted (h, q) -Bernoulli numbers and polynomials related to twisted (h, q) -zeta function and L -function, J. Math. Anal. Appl., 324(2006), 790-804.
- [24] Simsek, Y., On p -Adic Twisted q - L -Functions Related to Generalized Twisted Bernoulli Numbers, Russian J. Math. Phys., 13(3)(2006), 340-348.
- [25] Zill, D., and Cullen, M. R., Advanced Engineering Mathematics, Jones and Bartlett, 2005.

UNIVERSITY OF GAZIANTEP, FACULTY OF SCIENCE AND ARTS, DEPARTMENT OF MATHEMATICS,
27310 GAZIANTEP, TURKEY
E-mail address: mtsrkn@hotmail.com

UNIVERSITY OF GAZIANTEP, FACULTY OF SCIENCE AND ARTS, DEPARTMENT OF MATHEMATICS,
27310 GAZIANTEP, TURKEY
E-mail address: acikgoz@gantep.edu.tr